

OPPOSITIONAL DEFIANT DISORDER (ODD)

All children are oppositional from time to time, particularly when tired, hungry, stressed or upset. Oppositional behavior is often a common part of development for two to three year olds and early adolescents. However, openly uncooperative and hostile behavior becomes a serious concern when it becomes so frequent and consistent that it stands out when compared with other children of the same age and developmental level and when it affects the child's social, family and academic life.

In children with Oppositional Defiant Disorder (ODD) there is an ongoing pattern of uncooperative, defiant, and hostile behavior toward authority figures that seriously interferes with the youngster's day to day functioning.

Symptoms of ODD may include:

- ✓ **Frequent temper tantrums**
- ✓ **Excessive arguing with adults**
- ✓ **Active defiance and refusal to comply with adult requests and rules**
- ✓ **Deliberate attempts to annoy and upset people**
- ✓ **Blaming others for his or her mistakes and misbehavior**
- ✓ **Often being touchy or easily annoyed by others**
- ✓ **Frequent anger and resentment**
- ✓ **Mean and hateful talking when upset**
- ✓ **Seeking revenge**

Five to fifteen percent of school age children are reported to have ODD. It is important to look at other disorders that might be present such as ADHD (Attention Deficit Hyperactive Disorder), The Bi-Polar Child, depression, learning disabilities, and anxiety disorders (all of which we have strategies and remedies for).

It will be difficult to improve the symptoms of ODD without treating the co-existing disorder(s). A child with ODD can be challenging for the parents not only to cope with but very often the parents feel guilty. It is important to stress that this is a physiological disorder. Once again we can trace the behavioral symptomology to physiological imbalances rooting the issues. We have put together a summary of remedies to treat the vibration of the physiological pathogens to eliminate the behavioral symptoms, which make up "Oppositional Defiant Disorder".

Possible Complications

Be sure to test for the following issues

| | |
|-------------------------|---------------------------------|
| ADD | See ADD strategy |
| ADHD | See ADD strategy |
| CHILD BI-POLAR | See The Bi-Polar Child strategy |
| Vaccinations | See Vaccinations strategy |
| PENIERGOT | 1-2 |
| PENITUBERCULOSIS | 1-2 |

Grownups can have this condition too. They are always protesting, have trouble with authority figures, possibly paranoid, have trouble holding down a steady job and have difficulty in relationships.

In 2006 we formulated two remedies similar to ODD that affected attention and concentration. The symptoms are similar to many childhood issues listed in "Kid Stuff".

Phenyl Phrenosine Glucogenosis is similar to CDG, Congenital Disorder of Glycation (also called B GALTI deficiency). The first signs noted are :(1) Delayed speech development (2) lack of social communication (3) repetitive behavior (4) adapts to change poorly (5) "overloads" at social events (6) learning skills are diminished (7) low taste and smell ability.

Phenyl Hypo-Glycogenosis causes mood swings around digestion and after exercise. Personality changes from worthless to arrogant self-righteousness, all with strong doses of defiance. Starch is the largest cause of mental swings. In both childhood and adulthood concentration is difficult, mental frustration is high and muscle strength can vary wildly.

SUMMARY OF REMEDIES
With range of mega bottles needed
OPPOSITIONAL DEFIANT DISORDER 1-4